

AD-A171 868


AN OSTEOINDUCTIVE POLYMER COMPOSITE FOR CRANIAL AND
MAXILLOFACIAL BONE REPAIR (U) ARMY INST OF DENTAL
RESEARCH WASHINGTON DC J P SCHMITZ ET AL. 1985

1/1

UNCLASSIFIED

F/G 6/12

NL




END

DATE

FILED

10-86

17



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A171 868

(12)

AN OSTEOINDUCTIVE POLYMER COMPOSITE FOR
CRANIAL AND MAXILLOFACIAL BONE REPAIR

BY

John P. Schmitz D.D.S.

and

Jeffrey O. Hollinger D.D.S., PhD.

*U.S. Army Institute of Dental Research

Walter Reed Army Medical Center

Washington D.C. 20307-5300

DTIC
ELE
SEP 17 1986
A

* Address for reprints

Running title: Osteoinductive Polymer Composite

Key words: Bone Repair, Polymers, Osteoinduction

DTIC FILE COPY

This document has been approved
for public release and its
distribution is unlimited.

86 9 15 210

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO. <i>A171868</i>	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) An Osteoinductive Polymer Composite For Clinical and Maxillofacial Bone Repair.		5. TYPE OF REPORT & PERIOD COVERED
7. AUTHOR(s) Schmitz, J.P. Hollinger, J.O.		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS USAIDR WRAMC, Washington, DC 20307-5300		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE
		13. NUMBER OF PAGES
		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved For Public Release: Distribution is Unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) DISTRIBUTION STATEMENT A: Approved For Public Release: Distribution is Unlimited.		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Bone Repair, Polymers, Osteoinduction		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) See next page.		

20. ABSTRACT

A biodegradable copolymer of polylactide:polyglycolide (PLA:PGA) was combined with allogeneic decalcified freeze-dried bone (DFDB) and implanted into 15 mm diameter defects in the calvaria of 26 New Zealand White rabbits. Similar defects were created in the calvaria of another 26 rabbits. These animals served as controls and did not receive copolymer implants. Upon sacrifice, both the implants and the controls were evaluated clinically, radiographically, and histomorphometrically using a Zeiss Image Analysis System (Osteoplan[™] version 4.1). Both controls and implants were evaluated in groups at 4, 8, 12, 16, 20, and 24 weeks. When compared with the control defects, the copolymer:DFDB composite implants displayed a significantly greater volume of trabecular bone ($p < 0.025$). Two of the 15 mm diameter defects completely healed at 8 weeks. No adverse host tissue responses were observed histologically.

INSTRUCTIONS FOR PREPARATION OF REPORT DOCUMENTATION PAGE

RESPONSIBILITY. The controlling DoD office will be responsible for completion of the Report Documentation Page, DD Form 1473, in all technical reports prepared by or for DoD organizations.

CLASSIFICATION. Since this Report Documentation Page, DD Form 1473, is used in preparing announcements, bibliographies, and data banks, it should be unclassified if possible. If a classification is required, identify the classified items on the page by the appropriate symbol.

COMPLETION GUIDE

General. Make Blocks 1, 4, 5, 6, 7, 11, 13, 15, and 16 agree with the corresponding information on the report cover. Leave Blocks 2 and 3 blank.

Block 1. Report Number. Enter the unique alphanumeric report number shown on the cover.

Block 2. Government Accession No. Leave Blank. This space is for use by the Defense Documentation Center.

Block 3. Recipient's Catalog Number. Leave blank. This space is for the use of the report recipient to assist in future retrieval of the document.

Block 4. Title and Subtitle. Enter the title in all capital letters exactly as it appears on the publication. Titles should be unclassified whenever possible. Write out the English equivalent for Greek letters and mathematical symbols in the title (see "Abstracting Scientific and Technical Reports of Defense-sponsored RDT/E," AD-667 000). If the report has a subtitle, this subtitle should follow the main title, be separated by a comma or semicolon if appropriate, and be initially capitalized. If a publication has a title in a foreign language, translate the title into English and follow the English translation with the title in the original language. Make every effort to simplify the title before publication.

Block 5. Type of Report and Period Covered. Indicate here whether report is interim, final, etc., and, if applicable, inclusive dates of period covered, such as the life of a contract covered in a final contractor report.

Block 6. Performing Organization Report Number. Only numbers other than the official report number shown in Block 1, such as series numbers for in-house reports or a contractor/grantee number assigned by him, will be placed in this space. If no such numbers are used, leave this space blank.

Block 7. Author(s). Include corresponding information from the report cover. Give the name(s) of the author(s) in conventional order (for example, John R. Doe or, if author prefers, J. Robert Doe). In addition, list the affiliation of an author if it differs from that of the performing organization.

Block 8. Contract or Grant Number(s). For a contractor or grantee report, enter the complete contract or grant number(s) under which the work reported was accomplished. Leave blank in in-house reports.

Block 9. Performing Organization Name and Address. For in-house reports enter the name and address, including office symbol, of the performing activity. For contractor or grantee reports enter the name and address of the contractor or grantee who prepared the report and identify the appropriate corporate division, school, laboratory, etc., of the author. List city, state, and ZIP Code.

Block 10. Program Element, Project, Task Area, and Work Unit Numbers. Enter here the number code from the applicable Department of Defense form, such as the DD Form 1498, "Research and Technology Work Unit Summary" or the DD Form 1634, "Research and Development Planning Summary," which identifies the program element, project, task area, and work unit or equivalent under which the work was authorized.

Block 11. Controlling Office Name and Address. Enter the full, official name and address, including office symbol, of the controlling office. (Equates to funding/sponsoring agency. For definition see DoD Directive 5200.20, "Distribution Statements on Technical Documents.")

Block 12. Report Date. Enter here the day, month, and year or month and year as shown on the cover.

Block 13. Number of Pages. Enter the total number of pages.

Block 14. Monitoring Agency Name and Address (if different from Controlling Office). For use when the controlling or funding office does not directly administer a project, contract, or grant, but delegates the administrative responsibility to another organization.

Blocks 15 & 15a. Security Classification of the Report: Declassification/Downgrading Schedule of the Report. Enter in 15 the highest classification of the report. If appropriate, enter in 15a the declassification/downgrading schedule of the report, using the abbreviations for declassification/downgrading schedules listed in paragraph 4-207 of DoD 5200.1-R.

Block 16. Distribution Statement of the Report. Insert here the applicable distribution statement of the report from DoD Directive 5200.20, "Distribution Statements on Technical Documents."

Block 17. Distribution Statement (of the abstract entered in Block 20, if different from the distribution statement of the report). Insert here the applicable distribution statement of the abstract from DoD Directive 5200.20, "Distribution Statements on Technical Documents."

Block 18. Supplementary Notes. Enter information not included elsewhere but useful, such as: Prepared in cooperation with . . . Translation of (or by) . . . Presented at conference of . . . To be published in . . .

Block 19. Key Words. Select terms or short phrases that identify the principal subjects covered in the report, and are sufficiently specific and precise to be used as index entries for cataloging, conforming to standard terminology. The DoD "Thesaurus of Engineering and Scientific Terms" (TEST), AD-672 000, can be helpful.

Block 20. Abstract. The abstract should be a brief (not to exceed 200 words) factual summary of the most significant information contained in the report. If possible, the abstract of a classified report should be unclassified and the abstract to an unclassified report should consist of publicly-releasable information. If the report contains a significant bibliography or literature survey, mention it here. For information on preparing abstracts see "Abstracting Scientific and Technical Reports of Defense-Sponsored RDT&E," AD-667 000.

ABSTRACT

A biodegradable copolymer of polylactide:polyglycolide (PLA:PGA) was combined with allogeneic decalcified freeze-dried bone (DFDB) and implanted into 15 mm diameter defects in the calvaria of 26 New Zealand White rabbits. Similar defects were created in the calvaria of another 26 rabbits. These animals served as controls and did not receive copolymer implants. Upon sacrifice, both the implants and the controls were evaluated clinically, radiographically, and histomorphometrically using a Zeiss Image Analysis System, (OsteoplanTM version 4.1). Both controls and implants were evaluated in groups at 4, 8, 12, 16, 20, and 24 weeks. When compared with the control defects, the copolymer:DFDB composite implants displayed a significantly greater volume of trabecular bone ($p < 0.025$). Two of the 15 mm diameter defects completely healed at 8 weeks. No adverse host tissue responses were observed histologically.



DATE	
BY	
DISTRIBUTION/	
AVAILABILITY CODE	
Dist	Avail and/or Special
A-1	

INTRODUCTION

Avulsive maxillofacial wounds present challenging problems in reconstructive surgical management. These wounds often present as irregularly-shaped discontinuity defects that require some form of bone graft to restore continuity. Furthermore, defects in the maxillofacial region often occur on broad, flat regions of the skull and face where soft tissue support and facial esthetics are important requirements. A desirable repair material should be osteogenic, easily adaptable at the time of surgery, and rigid enough to support the muscles and soft tissues of the face.

For the repair of large maxillofacial defects, preferences are given to alloplastic materials, rib and iliac grafts, or calvarial grafts. Bone grafts present special problems including the need for a separate harvesting procedure, the potential limitation of available donor bone, and post-graft resorption. The advantages and disadvantages of autogenous grafts versus alloplastics have been reviewed by Wolfe[33].

The alpha-hydroxy polyesters, polylactide and polyglycolide, have been investigated for use as implant materials in the repair of a variety of soft tissue and osseous wounds. Implanted homopolymers and copolymers of polylactide and polyglycolide have been used in a supportive role as suture material, orbital floor replacements, dressings to facilitate healing in tooth extraction sites, and as biodegradable bone plates and screws[2, 3, 4, 5, 7, 23].

In the early 1960's, Urist discovered that consistent

osteoiduction by demineralized bone matrix could be achieved by the control of time, temperature, and molar concentration of HCl[29]. Allogeneic decalcified bone implants subsequently, have been used clinically for the correction of craniomaxillofacial deformities and in the treatment of jaw defects [8, 15, 19]. It was the purpose of this study to examine the osteogenic potential of a biodegradable copolymer(PLA:PGA) combined with allogeneic decalcified freeze-dried bone for craniofacial wound repair.

MATERIALS AND METHODS

Implant Preparation

A composite alloimplant was prepared by combining 50:50 poly(DL-lactide-co-glycolide) (PLA:PGA) with an inherent viscosity of 0.92 with allogeneic decalcified freeze-dried bone (DFDB). Diaphyseal segments of long bones were removed from donor animals (New Zealand White rabbits) and pulverized in the cold (8° C) to a particle size of 150-590 micrometers. The particles were defatted in 100% ethanol for four hours, decalcified for 24 hrs in 0.6 N HCl at 4° C, rinsed in 0.1M PBS and distilled water, and lyophilized. PLA:PGA copolymer (1 gm.) was solubilized in chloroform, precipitated with methanol and combined with DFDB (500 mg.). The doughy composite was forced into Teflon^R molds and heat cured for 24 hrs. at 45-48° C. After sterilization with ethylene oxide at room temperature, the implants were lyophilized for 84 hours at 50 millitorr to remove ethylene oxide residuals. Scanning electron micrographs of the cured implants revealed a polymer lattice interspersed with bone

matrix particles (Fig.1).

Surgery

Fifty-two, adult, male New Zealand white rabbits were randomly selected and conditioned for two weeks prior to the start of the experiment. Adult status was documented by radiographic confirmation of epiphyseal plate closure. The animals were anesthetized using a xylazine/ketamine technique supplemented with 1.8 ml of 2% lidocaine hydrochloride with 1:100,000 epinephrine. The scalp hair was removed with a depilating agent and the area over the calvaria was scrubbed with ethyl alcohol and povidone iodine for five minutes. Before surgery, 150,000 Units of Flocillin^R were administered intramuscularly in the left hind leg. Following attainment of a suitable level of anesthesia, a semi-lunar incision was made in the midline from the superior sagittal crest to the middle of the nasal bone. The soft tissues were gently reflected laterally and a 15 mm diameter craniotomy was created using a trephine in a slow-speed dental handpiece (Fig. 2a, 2b). Copious irrigation with normal saline was used throughout the procedure. Care was taken during the entire process to avoid perforating the dura or the superior sagittal sinus. The craniotomy was then irrigated with normal saline. Twenty-six of the animals served as controls while the other twenty-six animals received circular polymer composite implants (Fig. 3). The periosteum was closed over all the defects and implants and the tissues were closed in layers with 3-0 polyglycolic acid suture. At 4, 8, 12, 16, 20, and 24 weeks animals were

ethanatized with an overdose of sodium pentothal, USP. The defects and implants were retrieved along with surrounding host bone and immediately fixed in 70% ethanol. Specimens were removed from the fixative, placed dura-side down on Kodak Ultraspeed Dental X-ray film and radiographed at 90 KvP, 10 Ma, 0.4 seconds using a long-cone technique. Each radiograph was developed in an automatic x-ray processor and printed using a standard enlargement magnification. Specimens were embedded in polymethylmethacrylate, sectioned at 6 micrometers and stained with a modified Masson-Goldner trichrome stain. Specimens were analyzed using a Zeiss Image Analysis System (OsteoplanTM version 4.1). A random sampling of six fields for each specimen was measured. The bony fill per specimen was assessed by quantitating total trabecular bony volume (inclusive of calcified trabeculae and osteoid).

RESULTS

Gross Examination

Control sites

The control defects attempted to heal in a centripetal fashion. Although none of the 15 mm diameter defects healed in 24 weeks, healing was characterized by varying amounts of fibrous tissue repair (Fig. 4).

Implant sites

4 weeks: Extensive fibrous encapsulation of the implants was noted without significant degradation of the polymer (Fig. 5). The implants appeared to have swollen to the point of protruding out of the defect. The wound margins could

still be identified.

8 to 16 weeks: Varying amounts of osteosclerotic rimming were present at the wound margins with a reddish-brown soft tissue present within the center of the defect (Fig. 6). In many specimens the central area of the defect still contained areas of residual polymer.

20 to 24 weeks: Organized soft tissue elements and residual polymer were apparent. In implant sites which did not heal, dark red soft tissue was present in the center of the wound.

The implants were well tolerated by all the experimental animals. No adverse soft tissue reactions developed in any of the surgical sites. At necropsy, two of the implant-treated wounds healed completely at eight weeks. The copolymer implants were not completely degraded in all instances at 24 weeks, however, this did not appear to interfere with bone formation.

Radiographic Examination

Control sites

Control wounds demonstrated a centripetal pattern of bony repair characterized by finger-like extensions and occasional islands of osseous repair. None of the control wounds displayed radiographic evidence of osseous bridging at 24 weeks (Fig. 7).

Implant sites

The implants also demonstrated a centripetal pattern of bony repair. Two of the implants showed trabecular patterns indicative of complete repair at 8 weeks (Fig. 8). Coalescing

bony elements were present as irregular fingers and isolated areas of speckled radiopacities.

Histologic Examination

Control sites

4 weeks: Isolated islands of trabecular bone were present within the defect surrounded by fibrous connective tissue. Wound margins were not eburnated at this time.

8 to 12 weeks: Isolated areas of reparative elements were still present with apparent eburnation of the bony margins. The osseous islands appeared to increase in size with time. Many of these islands contained hematopoietic elements.

16 to 24 weeks: Islands of reparative elements became more numerous and incomplete attempts at osseous bridging were visible (Fig. 9).

Implant sites

4 weeks: Isolated islands of bony trabeculae were present throughout the defects. In the healing defects, widespread tessellation and rimming of osteoblasts was present (Fig. 10a, 10b). A minimal inflammatory cell infiltrate was present in areas between the demineralized bone particles.

8 to 12 weeks: Residual polymer was still present in the experimental defects. In three specimens, complete osseous bridging was observed (Fig. 11).

16 to 24 weeks: In areas devoid of healing, bone particles were occasionally present and appeared to be enveloped by fibrous connective tissue. Large islands of coalescing trabeculae were observed with areas of residual demineralized

bone surrounded by a prominent round cell infiltrate. Residual copolymer was present in many implant-treated wounds at 24 weeks.

Histomorphometric Analysis

The mean pooled trabecular bony volume from the experimental and control data was analyzed using an unpaired Student's t-test (Table I). There was a significant difference in the pooled mean trabecular bony volume between the implant and the control sites ($p < 0.025$).

DISCUSSION

Biodegradable polymers previously have been investigated for use in soft and hard tissue repair, in the internal fixation of fractures, and as intraosseous bone repair materials. A promising use for these polymers has been as carriers for osteogenic agents.

Nelson et al. were the first to assess the osteogenic potential of PLA/PGA copolymer implants in the repair of bony wounds [21]. The copolymer implants resulted in gradual healing of the bony wounds from the peripheries which progressed centrally. The implants were extremely tissue tolerant with little inflammatory or foreign body reaction. Olson et al. compared the tissue response of PLA with that of Gelfoam^R and Surgicel^R in healing extraction sites [23]. PLA was observed in the wounds after three months although its retention did not appear to interfere with new bone formation. PLA demonstrated less inflammatory reaction than either Gelfoam^R or Surgicel^R. Brekke et al. investigated the influence of PLA mesh on the incidence of localized osteitis [2]. They concluded that the use of PLA mesh substantially reduced the incidence of mandibular third molar "wound failure" and that the PLA demonstrated a hemostatic effect. Hollinger evaluated the osteogenic potential of PLA:PGA copolymer implants in osseous wounds [11]. He showed that although the implant material was still present at 42 days, it displayed an accelerated rate of healing as compared to control bony wounds. Hollinger was the first to use PLA:PGA as a carrier for a

calcification initiator, diposphoinositide-lysozyme (DPI-L) [12]. When DPI-L was combined with a PLA-PGA copolymer for the repair of endochondral wounds in rats and in the repair of mandibular discontinuity defects in dogs, the implants displayed an accelerated rate of repair as compared to the control wounds [12, 13]. Higashi et al. used PLA-hydroxyapatite (PLA-HA) composites to repair endochondral wounds in rats [10]. The PLA was observed to be rapidly resorbed and replaced by new bone which formed in direct contact with the HA.

Biocompatibility

There are no reports in the literature describing adverse systemic responses to polyester sutures or bone repair materials. Furthermore, the high degree of biocompatibility of PLA:PGA copolymers as bone implants has been described [1, 11, 18]. This was further supported by the high degree of tissue tolerance seen with the demineralized bone/copolymer composite. A PLA-HA composite was shown to become drastically acidic in vitro, reaching a pH of 3.4 in one week and remaining at that level for five weeks [10]. This localized pH drop may account for the lack of consistent healing pattern observed in many of the copolymer-DFDB composite implants.

Degradation

The process by which the alpha-hydroxy polyesters biodegrade is principally by non-specific hydrolytic scission [18]. Lactic acid, which is generated when PLA degrades, becomes incorporated into the tricarboxylic acid cycle, and is

excreted by the lungs as carbon dioxide and water [1]. PGA, on the other hand, is broken down hydrolytically as well as by nonspecific esterases and carboxypeptidases [32]. The resultant glycolic acid monomers are either excreted in the urine or enter the tricarboxylic acid cycle.

The rate at which the PLA:PGA implants degrade is dependent on the following six factors:

1. Molar ratio of the constituents.
2. Porosity of the implants.
3. Vascularity of the recipient site.
4. Degree of crystallinity of the constituents.
5. Average molecular weight.
6. Stress at the recipient site.

In general, PLA:PGA copolymers with a greater molar ratio of PLA tend to degrade more slowly than those with a greater amount of PGA [6, 18]. A 50:50 PLA:PGA molar ratio was chosen for this composite because its half-life appeared to be commensurate with normal fracture repair (four to six weeks) [18].

The total surface area of the polymer available also appears to influence the rate of degradation. Porous implants will degrade more rapidly than dense glassy implants. During fabrication of the implants used in this study, the control of porosity was possible only to a limited extent. An overriding concern was the fabrication of a rigid implant which could be carved and contoured at the time of surgery.

The degradation rate is also governed by the vascularity

of the recipient site. Implants placed in the mandible appear to degrade more rapidly than implants placed in the calvaria [14].

The degree of polymer crystallinity significantly effects the rate of water sorption. Consequently, the L(-) form degrades more slowly than the D,L form [16, 17]. Additionally, the alpha-hydroxy copolymers are less crystalline than their constituent homopolymers and will degrade more slowly [18]. A 50:50 poly(D,L-lactide-co-glycolide) implant was chosen because of its previous history of predictable degradation in endochondral wounds [11].

An important factor governing degradation is the concept of average molecular weight which is often described in terms of a polymer's inherent viscosity. Polymers that are highly viscous (high average molecular weight) will undergo slower biodegradation than those with a lower molecular weight and lower viscosity [22]. The presence of residual polymer at 24 weeks may be attributed to its viscosity in a relatively avascular site (e.g. calvaria). Previous studies with this same copolymer in the repair of mandibular discontinuity defects showed complete biodegradation by 8-12 weeks [H&S in press]. The flexing of the mandible which occurs in long-span discontinuity defects may be a factor in accelerating polymer degradation [14].

Mechanism of Action

Allogeneic decalcified freeze-dried bone is postulated to provide a substratum for the differentiation of perivascular

mesenchymal cells into chondrocytes and ultimately into osteoblasts. This process is known as osteoinduction. Urist has proposed that the inductor was a protein, specifically, bone morphogenetic protein (BMP) [30]. Reddi and coworkers, on the other hand, have proposed surface charge and geometry as well as inductor proteins as the putative factors in the osteoinduction process [24, 25, 27].

Following the heterotopic implantation of decalcified bone matrix, a sequential development of endochondral bone is initiated. Initially, the implant forms a plaque consisting of a conglomerate of decalcified matrix, fibrin, and neutrophils. Limited proteolysis is thought to cause the release of chemotactic factors and other matrix proteins into the surrounding milieu. The arrival of perivascular mesenchymal cells by chemotaxis is followed by their attachment to the bone matrix. The linear array of a random copolymer such as 50:50 PLA:PGA, which remotely resembles the linear array of the collagen molecule, may play a crucial role in the anchorage dependency of cells [28]. This anchorage dependency is essential in the early phase of the osteoinduction cascade. Cell attachment to bone matrix, thought to be promoted by fibronectin and related cell adhesive proteins, helps to bring the putative inductor in contact with focal cell surface receptors. The subsequent release of mitogenic factors from the matrix (hSGF) helps to promote growth and proliferation of cells [31]. These cells differentiate into chondrocytes and ultimately osteoblasts. Osteoblasts are subsequently seen to

be anchored to the periphery of the decalcified bone particles and to elaborate osteoid in a vectorial fashion.

Particulate cancellous marrow bone grafts presently are considered the material of choice for maxillofacial bone grafts. These grafts are used primarily to stabilize mobile skeletal segments, augment skeletal contour, and to construct new skeletal contour. However, the resorption of these grafts has been estimated to be in the range of 30-70% of graft bulk [19]. Demineralized bone powders, on the other hand, do not undergo resorption during bone induction and have complication rates no greater than that of conventional bone grafts. While impressive results have been documented clinically [8, 9, 15, 19, 20], Mulliken et al. [20] noted that "Bone powder fails to provide immediate stability and structure needed in many types of skeletal reconstruction and construction." In this regard, PLA:PGA copolymers may be a vehicle to provide the immediate stability and structure conducive to repair of the craniofacial skeleton.

REFERENCES

1. Brady, J.M., Cutright, D.E., Miller, R.A., Battistone, G.C., and Hunsuck, E.E. Resorption rate, route of elimination, and ultrastructure of the implant site of polylactic acid in the abdominal wall of the rat. *Oral Surg.* 7:155, 1973.
2. Brekke, J.H., Olson, R.A.J., Scully, J.R., and Osbon, D.B. Influence of polylactic acid mesh on the incidence of localized osteitis. *Oral Surg.* 56:240, 1983.
3. Christel, P., Chabot, F., Leray, J.L., Morin, C., and Mort, M. Biodegradable composites for internal fixation. In G.D. Winter, D.F. Giobbone, and H. Plena (Eds.), *Biomaterials*. New York: John Wiley and Sons, 1982. p.271.
4. Cutright, E.E. and Beasley, J.D. Fracture reduction using a biodegradable material, polylactic acid. *J. Oral Surg.* 29:393, 1971.
5. Cutright, D.E. and Hunsuck, E.E. The repair of fractures of the orbital floor using biodegradable polylactic acid. *Oral Surg.* 33:28, 1972.
6. Cutright, D.E., Perez, B., Beasley, J.D., Larson, W.L., and Posey, W.R. Degradation rates of polymers and copolymers of polylactic and polyglycolic acids. *Oral Surg.* 37:142, 1974.
7. Getter, L., Cutright, D.E., Bhaskar, S.N., and Augsburg, J.K. A biodegradable intraosseous appliance in the treatment of mandibular fractures. *J. Oral Surg.* 30:344, 1972.
8. Glowacki, J., Kaban, L.B., Murray, J.E., Folkman, J., and Mulliken, J.B. Application of the biological principle of induced osteogenesis for craniofacial defects. *Lancet* 1:959, 1981.
9. Glowacki, J. and Mulliken, J.B. Demineralized bone implants. *Clin. Plast. Surg.* 12:233, 1985.
10. Higashi, S., Yamamuro, T., Nakamura, T., Ikada, Y., Hyon, S.H., and Jamshidi, K. Polymer-hydroxyapatite composites for biodegradable bone fillers. *Biomaterials* 7:183, 1986.
11. Hollinger, J.D. Preliminary report on the osteogenic potential of a biodegradable copolymer of polylactide (PLA) and polyglycolide (PGA). *J. Biomed. Mater. Res.* 17:71, 1983.
12. Hollinger, J.D. Facilitation of osseous healing by a proteolipid copolymer material. Ph.D. Dissertation, University of Maryland. 1983.

13. Hollinger, J.O. and Schmitz, J.P. Restoration of bone discontinuity defects in dogs using a biodegradable implant. J. Oral Maxillofac. Surg. In press.
14. Hollinger, J.O. and Schmitz, J.P. Unpublished data. 1986.
15. Kaban, L.B., Mulliken, J.B., and Glowacki, J. Treatment of jaw defects with demineralized bone implants. J. Oral Maxillofac. Surg. 40:623, 1982.
16. Kulkarni, R.K., Pani, K.C., Neuman, C., and Leonard, F. Polylactic acid for surgical implants. Arch. Surg. 93:839, 1966.
17. Kulkarni, R.K., Moore, E.G., Hegyeli, A.F., and Leonard, F. Biodegradable poly(lactic acid) polymers. J. Biomed. Mater. Res. 5:169, 1971.
18. Miller, R.A., Brady, J.M., and Cutright, D.E. Degradation rates of oral resorbable implants (polylactates and polyglycolates): rate modification with changes in PLA/PGA copolymer ratios. J. Biomed. Mater. Res. 11:711, 1977.
19. Mulliken, J.B., Glowacki, J., Kaban, L.B., and Murray, J.E. Use of demineralized allogeneic bone implants for the correction of maxillocraniofacial deformities. Ann.Surg. 194:366, 1981.
20. Mulliken, J.B., Kaban, L.B., and Glowacki, J. Induced osteogenesis--the biological principle and clinical applications. J. Sur. Res. 37:487, 1984.
21. Nelson, J.F., Stanford, H.G., and Cutright, D.E. Evaluation and comparisons of biodegradable substances as osteogenic agents. Oral Surg. 43:836, 1977.
22. Odian, G. Principles of Polymerization, 2nd ed. New York:John Wiley and Sons, 1981.
23. Olson, R.A.J., Roberts, D.L., and Osbon, D.B. A comparative study of polylactic acid, gelfoam, and surgicel in healing extraction sites. Oral Surg. 53:441, 1982.
24. Reddi, A.H. and Huggins, C.B. Influence of geometry of transplanted tooth and bone on transformation of fibroblasts. Proc. Soc. Exp. Biol. Med. 143:643, 1973.
25. Reddi, A.H. and Huggins, C.B. Cyclic electrochemical inactivation and restoration of competence of bone matrix to transform fibroblasts. Proc. Natl. Acad. Sci. U.S.A. 71:1648, 1974.

26. Reddi, A.H. Implant-stimulated interface reactions during collagenous bone matrix-induced bone formation. *J. Biomed. Mater. Res.* 19:233, 1985.
27. Sampath, I.K. and Reddi, A.H. Dissociative extraction and reconstitution of extracellular matrix components involved in local bone differentiation. *Proc. Natl. Acad. Sci. U.S.A.* 78:7599, 1981.
28. Stoker, M., O'Neill, C., Berryman, S., and Waxman, U. Anchorage and growth regulation in normal and virus transformed cells. *Int. J. Cancer.* 3:683, 1968.
29. Urist, M.R. Bone: formation by autoinduction. *Science.* 150:893, 1965.
30. Urist, M.R., Mikulski, A., and Lietze, A. Solubilized and insolubilized bone morphogenetic protein. *Proc. Natl. Acad. Sci. U.S.A.* 76:1828, 1979.
31. Urist, M.R., DeLange, R.J., and Finerman, G.A.M. Bone cell differentiation and growth factors. *Science* 220:680, 1983.
32. Williams, D.F. and Mort, E. Enzyme-accelerated hydrolysis of polyglycolic acid. *J. Bioengin.* 1:23, 1977.
33. Wolfe, A.A. Autogenous bone grafts versus alloplastic materials in maxillofacial surgery. *Clin. Plast. Surg.*, 9:539, 1982.

TABLE 1. Trabecular Bony Volume of Composite
Implants Versus Controls

	Composite Implants	Controls
Six Field Volume (mm^3/cm^3)	1728.42	957.15
Std. Dev.	1614.44	869.15
n	26	26
p Value*	<0.025	

*Statistical data generated using an unpaired
Student's t-test

LEGENDS

Figure 1. Scanning electron micrograph of the PLA:PGA/DFDB composite. Particles of demineralized bone () can be seen to be invested within the copolymer matrix surrounded by open voids up to 200 micrometers wide (original magnification, X100.)

Figure 2a. Dried skull showing approximate size of the defect in relation to the overall calvaria.

Figure 2b. Surgical site following removal of the calvarial disc. Note the integrity of the dura and an intact superior sagittal sinus (arrow).

Figure 3. Polymer composite implant at the time of insertion demonstrating excellent adaptation to the bony wound margins.

Figure 4. 8 week control defect at retrieval. Fibrous repair tissue is present as a thin membrane over the defect. The wound margins are clearly demarcated (arrows).

Figure 5. 4 week implant at retrieval. Implant is found to have swollen within the defect with a fibrous tissue capsule present over the implant.

Figure 6. 8 week implant showing areas of osteosclerotic rimming (arrows).

Figure 7. Radiograph of a 24 week control defect with isolated bony islands present (arrows).

Figure 8. Radiograph of an 8 week implant showing trabecular patterns present throughout the defect.

Figure 9. Macrophotograph of a 24 week control defect. Note fibrous connective bridge.

Figure 10a. Tessellation and rimming of osteoblasts in a 4 week

specimen. Areas of amalgamated bone particles are present (arrows)(modified Masson-Goldner trichrome stain; original magnification,X64.)

Figure 10b. Anchorage of pre-osteoblasts to demineralized bone matrix in a 4 week implant. An active osteoblast may be seen anchored to the same particle(modified Masson-Goldner trichrome stain;original magnification,X100.)

Figure 11. Macrophotograph of a 20 week implant showing bony trabeculae continuous across the defect.

In conducting the research described in this report, the investigators adhered to the "Guide for the Care and Use of Laboratory Animals" as promulgated by the Committee on the Revision of the Guide for Laboratory Animal Facilities and Care of the Institute of Laboratory Animal Resources, National Research Council.

Commercial materials and equipment are identified in this report to specify the investigative procedure. Such identification does not imply recommendation or endorsement, or that the materials and equipment are necessarily the best available for the purpose. Furthermore, the opinions expressed herein are those of the authors and are not to be construed as those of the U.S. Army Medical Department.



10KV X100 100U 012 10263 IDR





Animal 53
VVR5 D-01-8
A1-41
13 Feb. 84



Animal

VVR5 D-01-84

B8-54

13 Feb. 84




Animal 50

VVR5 D-01-8.

B8-54

9 Apr. 84



Animal 46

VVR5 D-01-84

B4-42

9 Apr. 84




Animal 60

VVR5 D-01-84

A8-50

13 Mar.84





Animal 50

VVR5 D-01-84

B8-54

9 Apr. 84















DATA
FILM
O—